

Amphibia-Reptilia

Distribution and diversity of amphibians in Albania: new data and foundations of a comprehensive database --Manuscript Draft--

Manuscript Number:		
Full Title:	Distribution and diversity of amphibians in Albania: new data and foundations of a comprehensive database	
Short Title:	Amphibians in Albania	
Article Type:	Article	
Corresponding Author:	Márton Szabolcs HUNGARY	
Corresponding Author's Institution:		
First Author:	Márton Szabolcs	
Order of Authors:	Márton Szabolcs	
	Edvárd Mizsei	
	Daniel Jablonski	
	Balázs Vági	
	Béla Mester	
	Zsolt Végyári	
	Szabolcs Lengyel	
Manuscript Region of Origin:	ALBANIA	
Abstract:	<p>Albania is part of the Balkan Mediterranean biodiversity hotspot. Yet its herpetofauna is poorly known due to little scientific exploration during the long isolation of the country. To fill this gap, we constructed a georeferenced database on occurrences of all known amphibian species in the country based on records from published sources and on our data collected in expeditions to poorly known areas. Our database includes 1097 records of 16 species from between 1920 and 2017. We aggregated these records and data on altitude, climate, land cover diversity and distance from the sea in 10×10-km grid cells to analyse patterns in amphibian diversity. The mean number of species per cell was $1.8 \pm \text{S.E. } 0.11$ (maximum: 10 species). Sampling effort was uneven and sampling hotspots were mostly in popular natural heritage sites. Cells with high amphibian diversity were near Prokletije Mountains in the North-West, near Lura, Korab and Grammos Mountains, and Ohrid and Prespa Lakes in the East, and near Çikës Mountains and in coastal areas of Vlorë in the South-West. General linear models showed that the most important predictors of presence and diversity of amphibian species were land cover diversity and precipitation. Our study presents the largest database of amphibian occurrences in Albania to date. Our findings confirm the high richness and diversity of amphibians, mostly in areas with diverse land cover and high precipitation. Our study will be useful in biogeographic and ecological studies and for conservation purposes.</p>	
Keywords:	Balkan Peninsula; range; species richness; biogeography; BIOCLIM; GLMM	
Funding Information:	Országos Tudományos Kutatási Alapprogramok (K106133)	Mr. Márton Szabolcs Béla Mester Szabolcs Lengyel
	Agentúra na Podporu Výskumu a Vývoja (APVV-15-0147)	Daniel Jablonski
	Vedecká Grantová Agentúra MŠVVaŠ SR a SAV (SK) (1/0073/14)	Daniel Jablonski

	Societas Europaea Herpetologica (Mapping and Distributions 2015)	Mr. Márton Szabolcs Edvárd Mizsei Daniel Jablonski
	Balassi Intézet (B2/1CS/11521)	Mr. Márton Szabolcs
	Balassi Intézet (B2/1CS/19196)	Mr. Márton Szabolcs Edvárd Mizsei

Full title: Distribution and diversity of amphibians in Albania: new data and foundations
of a comprehensive database

Short title: Amphibians in Albania

Márton Szabolcs^{1,2,*}, Edvárd Mizsei³, Daniel Jablonski⁴, Balázs Vági³, Béla Mester^{1,2},
Zsolt Végyvári^{3,5}, Szabolcs Lengyel¹

¹Department of Tisza River Research, Danube Research Institute, Centre for Ecological
Research, Hungarian Academy of Sciences, Bem tér 18/c, 4026 Debrecen, Hungary

²PálJuhász-Nagy Doctoral School of Biology and Environmental Sciences, University
of Debrecen, Egyetem tér 1, 4032 Debrecen, Hungary

³Department of Evolutionary Zoology and Human Biology, University of Debrecen,
Egyetem tér 1, 4032 Debrecen, Hungary

⁴ Department of Zoology, Comenius University in Bratislava, Mlynská dolina,
Ilkovičova 6, 842 15 Bratislava, Slovakia

⁵ Department of Conservation Zoology, University of Debrecen – Hortobágy National
Park Directorate, H-4024 Debrecen, Sumen u. 2

*Corresponding author: szabolcs.marci@gmail.com

Article

Number of words in the manuscript: 7151

Number of words in the abstract: 239

Abstract: Albania is part of the Balkan Mediterranean biodiversity hotspot. Yet its herpetofauna is poorly known due to little scientific exploration during the long isolation of the country. To fill this gap, we constructed a georeferenced database on occurrences of all known amphibian species in the country based on records from published sources and on our data collected in expeditions to poorly known areas. Our database includes 1097 records of 16 species from between 1920 and 2017. We aggregated these records and data on altitude, climate, land cover diversity and distance from the sea in 10×10-km grid cells to analyse patterns in amphibian diversity. The mean number of species per cell was $1.8 \pm \text{S.E. } 0.11$ (maximum: 10 species). Sampling effort was uneven and sampling hotspots were mostly in popular natural heritage sites. Cells with high amphibian diversity were near Prokletije Mountains in the North-West, near Lura, Korab and Grammos Mountains, and Ohrid and Prespa Lakes in the East, and near Çikës Mountains and in coastal areas of Vlorë in the South-West. General linear models showed that the most important predictors of presence and diversity of amphibian species were land cover diversity and precipitation. Our study presents the largest database of amphibian occurrences in Albania to date. Our findings confirm the high richness and diversity of amphibians, mostly in areas with diverse land cover and high precipitation. Our study will be useful in biogeographic and ecological studies and for conservation purposes.

Key words: Balkan Peninsula, range, species richness, biogeography, BIOCLIM, GLMM

Introduction

Exploration and understanding the spatial distribution of biodiversity are one of the principal objectives of ecology (Gaston, 2000). Mapping species distributions and diversity also is a prime objective and tool in conservation (Pimm and Jenkins, 2005). However, the collection of records is often spatially biased, even in Europe.

Globally, amphibians are considered as one of the most threatened groups of animals (Gibbons et al., 2000, Alroy, 2015), where almost half of the species are declining (Stuart et al., 2004). The rapid decline of amphibians is explained by several factors such as fragmentation, degradation and complete loss of their habitats, global climate change, rapidly spreading diseases and synergies between these threats (Cushman, 2006; Sodhi et al., 2008). Rare species with restricted ranges and small populations are more likely to decline and to be affected by extinction risk (Harnik, Simpson and Payne, 2012). Moreover, some of amphibian species/endemic phylogenetic lineages are better adapted to past refugial regions and can survive better there than in current post-glacial ranges (Dufresnes and Perrin, 2015). Thus, information on species occurrences and their ranges are of key importance for conservation.

Albania is located in the western part of the Balkan Peninsula and is part of the Mediterranean hotspot of biodiversity (Griffiths, Kryštufek and Reed, 2004; Myers, 2000). The country covers 28,748 km² with an altitudinal range from sea level to 2764 meters. Collection of faunistic data on amphibians started in the early 20th century (Kopstein and Wettstein, 1920; Werner, 1920). This was followed by a long period when sampling lost its intensity until the next review on amphibians was published in the mid-nineties (Haxhiu, 1994). However, these data are more modest than those for

reptiles from the same period (see Haxhiu, 1998; Jablonski, 2011; Mizsei et al., in press). After amphibians came into the spotlight of conservation from the early 1990s due to their rapid decline and after the former isolationist political system ended in Albania in 1992, the number of records on amphibian species from Albania started to increase again in the early 21th century. However, many of these records remained unpublished in scientific papers.

Related mainly to the north-south orientation of Albania, the country covers the distribution of a wide range of amphibian species occurring in the Balkan Peninsula. Moreover, despite its small area, Albania has a rather high geomorphological variety and highly varied topography, for example, 70% of its terrain is mountainous (Fig 1). These conditions, along with a favourable Mediterranean climate led to the formation of a diverse pool of amphibian species (Pabijan et al., 2015). Landscape topography is primarily explained by orogenic processes affecting Albania and its surrounding areas as a consequence of the collision of the Adria microplate with the Eurasian plate. During the last stage of the Neotectonic Pliocene-Quaternary period, from the Middle Pleistocene to the present times, local episodes of subsidence in Albania induced the formation of graben lakes such as Shkodra, Ohrid and Prespa, and the development of Quaternary graben plains (Aliaj, Baldassare and Shkupi, 2001). This land evolution, in combination with the later relative stability of the Mediterranean climate resulting from little influence of Pleistocene glaciations, led to allopatric speciation and differentiation of amphibians. There are strong indications that some of the Miocene-Pliocene speciation centres or Pleistocene glacial refugia of amphibians (and other taxa with limited dispersal ability) were located inside or in close to the current territory of Albania (Médail and Diadema, 2009). The Western Balkan is home to two endemic

species of water frogs (*Pelophylax epeiroticus*, *P. shqipericus*), one endemic brown frog (*Rana graeca*), one endemic crested newt (*Triturus macedonicus*) and smooth newt (*Lissotriton graecus*) which are also part of the Albanian amphibian fauna (Sillero et al., 2014a; Pabijan et al., 2015; 2016). However, detailed data on the genetic diversity of amphibian species in Albania are still lacking. Current coastal and freshwater ecosystems include rivers, streams, lakes, swamps, estuaries, lagoons or drainage channels and all these represent suitable habitats for amphibians. To date, 16 species of amphibians have been detected in Albania (Haxhiu, 1994; Szabolcs and Mizsei, in press). Mainly due to the southern location and the landscape- and habitat-scale heterogeneity of Albania, this number of species is higher than in many other European countries covering a larger area (Haxhiu, 1994). Therefore, this region has central importance in understanding both the past and present patterns of amphibian diversity in Europe, which thus warrants a synthesis and an update of the current knowledge on amphibian species of the country.

The aims of our study were (i) to fill gaps in our knowledge on the distribution of amphibian species in Albania by collecting occurrence records from previous literature and supplementing them by our recently collected data into a single georeferenced database, (ii) to present up-to-date distribution maps for each species, and (iii) to analyse patterns in species diversity in order to find hotspots of amphibian diversity and to identify environmental factors explaining the distribution of amphibians in Albania. This study is complementary to our previous study on the distribution and diversity of reptiles in Albania (Mizsei et al., in press).

Materials and Methods

Data collection and processing

We used five sources of data to collate our database on the occurrences of amphibians in Albania. First, we collected records by searching the primary literature for studies and reports of amphibian species in Albania. Whenever it was possible, we georeferenced published maps in Quantum GIS 1.8.0 using the GDAL plugin (Bruno, 1989) or used the original field-collected coordinates (as in Bringsøe, 2011; Jablonski, 2011; Recuero et al., 2012; Pabijan et al., 2015; Szabolcs and Mizsei, in press). If maps or coordinates were not available (as in Kopstein and Wettstein, 1920; Werner, 1920; Frommhold, 1962; Schneider and Haxhiu, 1994; Haxhiu, 1994; 2000a; 2000b; 2000c; 2000d; Farkas and Búzás, 1997; Ragghianti et al., 1999; Haxhiu and Vrenozi, 2009; Shehu et al., 2009; Oruçi, 2010; Aliko et al., 2012; Denoël et al., 2012; Guignard et al., 2012; Aliko, Biba & Sula, 2013; Aliko, Qirjo & Nuna, 2014; Shkurti, 2013), we identified localities given in the studies using Google Earth 7.1.8, Google Maps (<http://maps.google.com>), the GeoNames database (<http://geonames.org>) and various other websites and blogs. Second, we processed records from the amphibian collection of the Hungarian Natural History Museum. Third, we added records from the Global Biodiversity Information Facility (GBIF, <http://gbif.org>, which includes records from several museums), iNaturalist (<http://inaturalist.org>) and TrekNature (<http://treknature.com>) databases with the permission of the data providers. Fourth, we obtained records from fellow scientists and citizen herpetologists with extensive knowledge of Albanian amphibians. The internet forum called Fieldherping.eu (<http://fieldherping.eu>) was a major source to contact these experts. Fifth, we added our own unpublished data collected in a total of 21 expeditions to Albania. Most of these expeditions were conducted as part of studies on the Greek Meadow Viper (*Vipera graeca*, Mizsei et al. 2016), but we also visited areas from where we found no information in the above four sources during these trips to search for amphibians. We stored all records in point shapefiles in a GIS database.

For species treatment, we used the most up-to-date nomenclature and taxonomy from Sillero et al. (2014a) and Speybroeck et al. (2016). Three species from the frog genus *Pelophylax* (*P. epeiroticus*, *P. ridibundus* and *P. shqipericus*) are difficult to identify based on external morphological characters, use similar habitats and are known to hybridise with each other. Although most of the *Pelophylax* species can be distinguished based acoustically (Schneider and Haxhiu, 1994, Lukanov, Tzankov & Simeonovska-Nikolova, 2015) we did not have this information in most cases, thus we merged these three species into

Pelophylax spp. to avoid the possibility of sampling bias to any of the three species (see e.g. Mester et al., 2015). According to recent taxonomical changes it seems that two species occur in Albania from the genus *Bufo* (Özdemir et al., 2014). Because we could not differentiate them in the field we merged them under the name *B. viridis/variabilis*. We used the name *Lissotriton graecus* instead of *L. vulgaris* because recent molecular analyses are indicating the species status of this Balkan lineage (Pabijan et al., 2016).

For spatial visualisation and analyses, we aggregated point records into a 10×10 km grid system provided by the European Environmental Agency (<http://eea.europa.eu/data-and-maps/data/eea-reference-grids>) in ETRS89 Lamberth Azimuthal Equal Area projection (EPSG: 3035). To identify the elevation of the localities in order to determine the altitudinal range of the species, we used the Shuttle Radar Topographic Mission (SRTM) 90-m Digital Elevation Database 4.1 (Jarvis et al., 2008). We also noted the year for every record.

Spatial analyses

Spatial autocorrelation among the records of the dataset and bias due to spatially uneven sampling are common biases in point occurrence data (Rocchini et al., 2011). We tested for spatial autocorrelation in the number of records per cell using the global Moran's I spatial statistic. This statistic tests the null hypothesis that the occurrence records are evenly distributed against the alternative hypothesis that the records are spatially either clustered ($Z > 0$) or dispersed ($Z < 0$). To analyse patterns in sampling bias, we used the Getis Ord G_i^* spatial statistic (Ord and Getis, 1995), which informs whether sampling effort is significantly lower (G_i^* score < -1.96 , coldspot of sampling) or higher (G_i^* score > 1.96 , hotspot of sampling) than expected by chance. We used ESRI ArcGIS 10.0 in these analyses.

We calculated Shannon diversity for each 10×10 km cell and then visualised the occurrences of the species within the cells. Additionally, we calculated the Extent of Occurrence (EOO) for each species as the number by fitting a Minimum Convex Polygon to their point records and then dissected it with the territory of Albania to obtain the Albanian range of each species.

Environmental data and linear modelling

We obtained information on several environmental variables to test which factors correlate with patterns in amphibian presence/absence and diversity (Table 1). First, we obtained data on 19 Bioclim variables from the WorldClim database (Hijmans et al., 2005). We then applied a principal component analysis to extract four principal components, which explained 99% of the total variance. Second, to characterise habitat diversity, we calculated the Shannon-diversity of CORINE Land Cover (250 m resolution; European Environmental Agency) in each cell using the LecoS 1.9.8 plugin in QGIS (Jung, 2012). Third, to characterise elevation and altitudinal variation within the cells, we calculated the mean and standard deviation (S.D.) of altitude within each 10×10 km cell based on grid values of the Shuttle Radar Topographic Mission (SRTM) 90-m Digital Elevation Database 4.1 (Jarvis et al., 2008) using Zonal Statistics in QGIS 2.12. Finally, we calculated the distance between the centroid of each cell and the closest point to the sea shore using the NNJoin 1.2.2 plugin in QGIS.

To evaluate the effects of environmental variables on amphibian presence/absence, we applied a model selection approach by fitting generalized linear mixed models (GLMM) with binomial error distribution (Pinheiro and Bates, 2000) using the lme4 package in the R environment (R Core Team, 2015). We ran models for all possible combinations of the environmental variables. To evaluate the relation between Shannon diversity and environmental predictors, we fitted a GLMM using the Markov chain Monte Carlo (MCMC) routine of the MCMCglmm package with its default parameters (Hadfield, 2010). To control for spatial autocorrelation, we specified cell ID as a random factor, and to control for sampling bias, we included GiZ scores as a random factor in both modelling approaches. To minimise the influence of phylogenetic relatedness of the species, we included species ID nested in order as an additional random factor in the GLMM for amphibian presence.

After model selection, we calculated the relative importance of environmental predictors using model-comparison techniques in an information-theoretic framework (Burnham and Anderson, 2002). In the first step, we obtained the values of Akaike's information criterion corrected (AICc) for small sample sizes, which is a metric of the trade-off between the goodness of fit of the model and its complexity, thus functioning as a measure of information entropy. Next, we assessed the corresponding Akaike weight of each model (ω) representing the relative likelihood of a model. In the third step, we selected models with substantial support: Akaike differences in the range 0-2 indicate substantial level of empirical support of a given model ($\Delta_i = \text{AIC}_i - \text{AIC}_{\min} < 2.0$) (Burnham and Anderson, 2002). We calculated model-averaged

parameter estimates (θ) and unconditional standard errors that controlled for model uncertainty (SEu; Burnham and Anderson, 2002) of each variable by the sums of their Akaike weights across all models with substantial support containing the given predictor.

For all analyses, we used the R 3.3.2 statistical computing environment (R Core Team, 2015). Model fitting and selection were performed applying the MuMIn package (Barton, 2011).

Results

Distributional evaluation

We collected a total of $N = 1097$ amphibian occurrence records. The earliest records were from 1920, and the rate of collection was slow until the mid-20th century (Fig. 2). After 1962, the number of records started to increase, with one large peak in 1994 (Haxhiu, 1994). Nearly half of the total number of occurrence records ($N = 555$) are published here for the first time, while the other half ($N = 542$) were published previously (Table 2, Fig. 3a). Of the unpublished records, we collected a total of $N = 482$ records during our expeditions, $N = 18$ from museum collection, $N = 8$ from internet sources and added $N = 47$ records from personal communication. The number of records ranged from three in the case of *Pelobates syriacus* to 339 in *Pelophylax* spp. For most species, half or nearly half of the records were unpublished, and we had at least one unpublished record for all species (except *P. syriacus*) (Table 2). In the database, *P. syriacus* was also the rarest species, present in only one grid cell, while *Pelophylax* spp. was the most widely distributed taxon, present in 181 grid cells. At least one species of amphibian occurred in 238 of the 349 grid cells covering Albania. Measurement of the EOO revealed that many species with only a few records had much larger possible range than expected. For instance, *Lissotriton graecus* was present in 47

or *Rana graeca* in 43 cells, however, their EOO was close to the total area of the country.

Amphibian diversity patterns

According to Global Moran's *I* spatial statistics, overall sampling effort in Albania was spatially clustered ($Z = 4.064$, $P < 0.0001$). Although Getis Ord G_i^* statistics did not reveal coldspots of sampling effort, sampling hotspots were found (Fig. 3b), mostly in the Prokletije Mountains (Mts.), the vicinity of Ohrid and Prespa Lakes, Pindos Mts., coastal regions near Vlorë and around Butrint Lake in the south (Fig. 3c).

The mean number of species per cell was $1.8 \pm (\text{S.E.}) 0.11$, with a maximum of $N = 10$ in two cells. Cells with high amphibian diversity occurred in Prokletije Mts., Lura and Korab Mts., vicinity of Ohrid and Prespa Lakes, Grammos Mts., Çikës Mts. and coastal regions near Vlorë (Fig. 3c). Distribution maps of amphibians of Albania are presented in the Supplementary material S1-S16.

Most amphibians had a large altitudinal range between sea level to above 1500 m a. s. l., with the exception of a few, mostly mountain-dwelling species (e. g. *Salamandra atra*, *Rana temporaria*) which had lower sample sizes and/or narrower ranges (Fig. 4). GLMM models showed that the most important predictors for amphibian presence and diversity were land cover diversity (CORINE DIV) and precipitation (BIO PC2), whereas temperature variation (BIO PC3) was important for amphibian presence only (Table 3). Each of these variables was part of at least one of the best models for presence and diversity (Table 4). Model-averaged parameter estimates suggested that CORINE DIV (land cover diversity), BIO PC2 (precipitation) and BIO PC3 (temperature variation) significantly influenced amphibian presence,

whereas diversity was influenced only by CORINE DIV (Table 5). The effect of CORINE DIV was positive for both presence and diversity, whereas that of BIO PC2 was negative for presence (Table 5, Fig. 5).

Discussion

Our study presents a spatially explicit database containing the largest amount of amphibian records from Albania to date and a fine-scale analysis of patterns in amphibian occurrence and diversity involving all known species in the country.

The relatively low species richness in many cells indicates that most of the country is still data deficient (Fig. 3b). Amphibian biodiversity hotspots were found where sampling effort was higher than average (i.e., in protected areas or popular tourist destinations such as Theth and Prespa National Parks, Butrint World Heritage Site). These results are similar to those of Cogălniceanu et al. (2013) from Romania, a country also characterized by uneven sampling mainly due to high altitudinal complexity and uneven road density. Because amphibian species richness per cell was relatively low in our study (c.f. mean number of species per cell: 1.68), rare species exerted a large impact on the designation of amphibian hotspots. Among them we found *Pelobates syriacus* which reaches its westernmost distribution in a single cell in the southeast (Szabolcs & Mizsei, in press), and montane species *Ichthyosaura alpestris*, *Rana temporaria* and *S. atra* with a relatively restricted altitudinal distribution (Table 1; Fig.4). These results imply that the low number of records for some rare species cannot be expanded by further sampling due to biogeographic constraints determining their range (Bruno, 1989). Although we found no evidence of sampling coldspots and the altitudinal distribution of records corresponded well with the frequency distribution of

altitudinal values of Albania (Fig. 4), it was clear from the generally low species richness that there is a need to add more data on the occurrences of species other than these rare specialists.

The temporal distribution of our records showed that data collection was almost halted during the Communist era (1946-1992), then restarted during the 1990s, and more recently it has yielded an unprecedented amount of records. This corresponds to the abandonment of the political and economic isolation of the previous regime, which also resulted in higher standards of living and a dynamically growing GDP (<http://imf.org>). However, economic development also leads to the abandonment of traditional land uses and an increasing rate of habitat alterations, both of which usually have a negative impact on amphibian diversity (Scribner et al., 2001; Hartel et al., 2009). Examples are shown by an increasing number of road constructions and large-scale hydropower projects for example in the valleys of Vjosë or Valbona rivers (Freyhof, 2010; <http://balkanrivers.net>), which may lead to the devastation of important wetland habitats (Cushman, 2006).

Our analysis of the data currently available showed that the diversity of land cover was the most important factor affecting both the occurrence and the diversity of amphibian species in Albania. These results agree well with those of several previous landscape-scale studies (e.g. Van Buskirk, 2005; Denoël and Ficetola, 2008; Hartel et al., 2009; Vági et al., 2013) and can be explained by two mutually non-exclusive hypotheses. First, most amphibians are characterised by a complex life cycle and they use different environments during their larval and adult life, thus, most species are expected to require complex habitats. Second, those species which use aquatic habitats all year round even as adults can mainly spread along aquatic habitats (Ficetola and De

Bernardi, 2004), thus their occurrence is primarily related to local hydrological factors rather than to climate or land use and they can occur under various climatic conditions and land use types. Indeed, in our study, the species with the largest amount of records were *Pelophylax* spp. and *Bombina variegata* (Table 2.). Both of these anurans are associated with freshwater habitats all year round (Arnold and Ovenden, 2002). The relatively large number of records for these two species can also be explained by their high detectability: these anurans are frequent in several types of waters due to their wide ecological tolerance, are often active in daylight, and calling males can be easily detected acoustically even in the hottest summer months due to their prolonged breeding season (Arnold and Ovenden, 2002). All these factors can lead to sampling bias in mapping surveys (Cogălniceanu et al., 2013).

In contrast, other amphibians are mostly terrestrial throughout the year, are often active only in rainy and moist weather, mainly at night, thus are more difficult to detect in the field. These terrestrial, but still widespread species can be characterized by a large, country-wide EOO often with a wide altitudinal range, even if they were detected only in a small fraction of cells. Species in this group included *Bufo bufo*, *Bufotes viridis/variabilis*, *Hyla arborea*, *Lissotriton graecus*, *Rana dalmatina* and *Triturus macedonicus* (Table 2.). Two other species, *Rana graeca* and *Salamandra salamandra* also showed large EOO because these species are mostly associated to mountain habitats, and can thus be widespread in Albania (Table 2.). Adequate sampling of these species requires surveys during the breeding season in spring, when most amphibians stay in and around water bodies all day. The detection probability of individuals in various life stages can be further increased by a combination of newt traps, dip-netting

and visual or acoustic surveys (Ficetola and De Bernardi, 2004; Van Buskirk, 2005; Mattfeldt, 2007; Vági et al., 2013; Mester et al., 2015).

We did not distinguish between the three *Pelophylax* species occurring in Albania, as they are hard to identify based only on morphological characters. *P. ridibundus* is capable of hybridisation with the two other species (Schneider and Haxhiu, 1994; Ragghianti et al., 1999; 2004) which also makes their identification difficult. *P. epeiroticus* is found between the extreme south-western Albania to southern Greece along the Ionian coast and is genetically related to *P. ridibundus* (Lymberakis et al., 2007). *P. shqipericus* lives in southern Montenegro from Lake Shkodra to Vlora Bay in western Albania along the Adriatic coast. This species is genetically related to *P. lessonae* (Ragghianti et al., 2004). *P. ridibundus* has a country-wide distribution and is common in the Balkans and Europe (Sillero et al., 2014a). The latter species may co-occur with the two former ones although we found no evidence for habitat overlap between the three species. Little is known about the ecology of the two Balkan species and their coexistence with *P. ridibundus*, therefore, further research is highly necessary as both are threatened with extinction. IUCN Red List category of *P. epeiroticus* is Vulnerable (Uzzell, Lymberakis & Haxhiu, 2009) and of *P. shqipericus* is Endangered (Uzzell and Crnobrnja-Isailović, 2009).

Besides habitat alteration (loss, fragmentation, degradation), other threats to amphibians include climate change and spread of diseases. Climate change can alter the amount and distribution of precipitation, which was among the most important factors governing amphibian occurrences (Table 5). Finally, the chytrid fungus *Batrachochytrium dendrobatidis* (Fisher, Garner and Walker, 2009) has been also detected in eight species of Albanian amphibians (Vojar et al., 2017), although we are

not aware about any outbreak of the disease chytridiomycosis in the East Mediterranean. In conclusion, a detailed assessment of the distribution of amphibian species and diversity and an evaluation of the efficiency of protected and properly managed areas is urgently needed in Albania. We hope that our work will be an important starting point toward this aim. To facilitate future work, the spatially explicit database and methodological approaches presented here provide important baseline information. Our results can be integrated to larger databases such as the NA2RE – New Atlas of Amphibians and Reptiles of Europe (Sillero et al., 2014a; 2014b; <http://na2re.ismai.pt>).

Acknowledgements

Many people contributed to this work as data providers namely W. Babik, B. Demovič, M. Durakic, F. Katallozi, M. Lemonnier-Darcemont, O. Suldovska, E. Sunje, G.J. Verspui, M. Vogrin, S. Voitel, F. Weber, M. Wolny and D. Zerzán. We are grateful to J. Vörös not only for cordially providing data of the Hungarian Natural History Museum, but for barcoding also a *Rana* metamorph. Financial support was provided by the CSEMETE Environmental and Nature Conservation Association; the Serpentes Foundation; the Department of Evolutionary Zoology, University of Debrecen; the Balassi Institute, Hungary (B2/1CS/11521, B2/1CS/19196); and a grant from OTKA (K106133). DJ was supported by the Slovak Research and Development Agency under the contract No. APVV-15-0147 and by grant of the Scientific Grant Agency of the Slovak Republic VEGA 1/0073/14. Fieldwork to collect data on amphibian occurrences was supported by a Grant in Herpetology – Mapping and Distributions 2015 from the Societas Europaea Herpetologica (SEH) to EM, MSz and DJ.

372

373 **References**

374 Aliaj, Sh., Baldassarre, G., Shkupi, D. (2001): Quaternary subsidence zones in Albania:
375 some case studies. Bull. Eng. Geol. Env. **59**: 313-318.

376 Aliko, V., Biba, A., Sula, E., Gjurgjaj, A. (2012): Effects of pollution on amphibian
377 blood parameters (Ranidae: *Rana balcanica* & *Rana lessonae*) from the
378 Albania's coastal zone. In: International Conference on Marine and Coastal
379 Ecosystems, p. 207-213. Anonymous (ed), Tirana.

380 Aliko, V., Biba, A., Sula, E. (2013): Clastogenic effects of Tirana lake water on
381 *Pelophylax kurtmuelleri* red blood cells. In: 3rd International Conference of
382 Ecosystems, p. 1-6. Dursun, S., Mankolli, H., Zuchetti, M., Vosniakos, F. K.
383 (eds), Tirana.

384 Aliko, V., Qirjo, M., Nuna, E. (2014): Haematological and morphological effects of
385 copper sulfate on the larval development of green toad, *Bufo viridis*. In: 4th
386 International Conference of Ecosystems, p. 736-743. Anonymous (ed), Tirana.

387 Alroy, J. (2015): Current extinction rates of reptiles and amphibians. Proceedings of the
388 National Academy of Sciences **112**: 13003-13008. Arnold, N., Ovenden, D.W.
389 (2002): Reptiles and Amphibians of Britain and Europe. London, Collins.

390 Bartón, K. (2016): Mumin: Multi-Model Inference. R package version 1.15.6.

391 Bringsøe, H. (2011): Possible circadian colour change in *Rana graeca* in Albania. Z.
392 Feldherpetol. **18**: 93-98.

393 Bruno, S. (1989): Introduction to a study of the herpetofauna of Albania. British
394 Herpetological Society Bulletin **29**.

395 Burnham, K. P., Anderson, D. R. (2002): Model selection and multimodel inference.
 396 New York, Springer-Verlag.

397 Cogălniceanu, D., Székely, P., Samoilă, C., Iosif, R., Tudor, M., Plăiașu, R., Stănescu,
 398 F., Rozyłowicz, L. (2013): Diversity and distribution of amphibians in Romania.
 399 ZooKeys **296**: 35-57.

400 Cushman, S. A. (2006): Effects of habitat loss and fragmentation on amphibians: A
 401 review and prospectus. Biol. Cons. **128**: 231-240.

402 Denoël, M., Duguet, R., Džukic, G., Kalezić, M., Mazzotti, S. (2001): Biogeography
 403 and ecology of paedomorphosis in *Triturus alpestris* (Amphibia, Caudata). J.
 404 Biogeogr. **28**: 1271-1280.

405 Denoël, M., Ficetola G. F., (2008): Conservation of newt guilds in an agricultural
 406 landscape in Belgium: the importance of aquatic and terrestrial habitats. Aquat.
 407 Conserv. **18**: 714-728.

408 Dufresnes, C., Perrin. N. (2015): Effect of biogeographic history on population
 409 vulnerability in European amphibians. Cons. Biol. **29**: 1235-1241.

410 Farkas, B., Buzás, B. (1997): Herpetologische Beobachtungen in den Nordalbanischen
 411 Alpen. Herpetofauna **19** (109): 10-18.

412 Ficetola, G.F., De Bernardi, F. (2004): Amphibians in a human-dominated landscape:
 413 the community structure is related to habitat features and isolation. Biol. Cons.
 414 **119**: 219-230.

415 Fisher, M. C., Garner, T. W. J., Walker S. F. (2009): Global emergence of
 416 *Batrachochytrium dendrobatidis* and amphibian chytridiomycosis in space, time,
 417 and host. Annu. Rev. Microbiol. **63**: 291-310.

418 Freyhof, J. (2010): Threatened mollusks and freshwater fish of the Balkans, Potential
 419 impacts of hydropower projects. Berlin, ECA Watch Austria & Euronatur, pp.
 420 81.

421 Frommhold, E. (1962): Herpetologische Studien in Albanien. Aquarium Terrarium **9**
 422 (12): 365-370.

423 Gaston, K. J. (2000): Global patterns I biodiversity. Nature **405**: 220-227.

424 Gibbons, J. W., Scott, D. E., Ryant, T. J., Buhlmann, K. A., Tuberville, T. D., Metts, B.
 425 S. (2000): The global decline of reptiles, Déjà Vu Amphibians. BioScience**50**:
 426 653-666.

427 Griffiths, H.I., Kryštufek, B., Reed, J.M. (2004): Balkan biodiversity: Pattern and
 428 Process in the European Hotspot. Dordrecht, Kluwer Academic Publisher.

429 Guignard, M., Büchi, L., Gétaz, M., Betto-Colliard, C., Stöck, M. (2012): Genome size
 430 rather than content might affect call properties in toads of three ploidy levels
 431 (Anura: Bufonidae: Bufo viridis subgroup). Biol. J. Linn. Soc. **105**: 584-590.

432 Hadfield, J. D. (2010): MCMC methods for multi-response generalized linear mixed
 433 models: the MCMCglmm R package. J. Stat. Softw. **33** (2): 1-22.

434 Harnik, P.G., Simpson, C., Payne, J.L. (2012): Long-term differences in extinction risk
 435 among the seven forms of rarity. Proc. R. Soc. Lond. B Biol. Sci. **279**: 4969-
 436 4976.

437 Hartel, T., Nemes, S., Cogălniceanu, D., Öllerer, K., Moga, C.I., Lesbarrères, D.,
 438 Demeter, L. (2009): Pond and landscape determinants of Rana dalmatina
 439 population sizes in a Romanian rural landscape. ActaOecol. **35**: 53-59.

440 Haxhiu, I. (1994): The herpetofauna of Albania, Amphibia: species composition,
 441 distribution, habitats. Zool. Jahrb. Syst. **121**: 321-334.

442 Haxhiu, I. (2000a): Herpetofauna in the Karaburun area. Scientific report.

443 Haxhiu, I. (2000b): Herpetofauna in the Llogara area. Scientific report.

444 Haxhiu, I. (2000c): Herpetofauna in the Narta area. Scientific report.

445 Haxhiu, I. (2000d): Herpetofauna in the Orikum area. Scientific report.

446 Haxhiu, I., Vrenozi, B. (2009): Species of amphibians and reptiles of lake Ohri with
 447 notes in their ecology. In: International Conference on Lakes and Nutrients,
 448 Pogradec (Albania), p. 382-387. Anonymus (ed), Pogradec.

449 Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones P.G., Jarvis, A. (2005): Very high
 450 resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**:
 451 1965-1978.

452 Jablonski, D. (2011): Reptiles and amphibians of Albania with new records and notes
 453 on occurrence and distribution. *Acta Soc. Zool. Bohem.* **75**: 223-238.

454 Jarvis, A., Reuter, H.I., Nelson, A., Guevera, E. (2008): Hole-filled SRTM for the globe
 455 v4.1 (<http://srtm.csi.cgiar.org>).

456 Jung, M. (2013): LecoS – A QGIS plugin for automated landscape ecology analysis.
 457 *PeerJ PrePrints* 1:e116v2.

458 Kopstein, F. Wettstein, O. (1920): Reptilien und Amphibien aus Albaniens. *Verh. Zool.*
 459 *Bot. Ges. Öst.* **70**: 387-457.

460 Lukanov S., Tzankov, N., Simeonovska-Nikolova D. (2013): A comparative study of
 461 the mating call of *Pelophylax ridibundus* and *Pelophylax kurtmuelleri* (Anura:
 462 Ranidae) from syntopic and allotopic populations. *Journal of Natural History* **49**:
 463 257-272.

464 Lymberakis, P., Poulakakis, N., Manthou, G., Tsigenopoulos, C. S., Magoulas, A.,
 465 Mylonas, M. (2007): Mitochondrial phylogeography of *Rana* (*Pelophylax*)

466 populations in the Eastern Mediterranean region. *Mol. Phylogenet. Evol.* **44**:
 467 115-125.

468 Mattfeldt, S. D., Grant E. H. C. (2007): Are two methods better than one? Area
 469 constrained transects and leaf litterbags for sampling stream salamanders.
 470 *Herpetological Review* **38**: 43-55.

471 Médail, F., Diadema, K. (2009): Glacial refugia influence plant diversity patterns in the
 472 Mediterranean Basin. *J. Biogeogr.* **36**:1333-1345.

473 Mester, B., Szalai, M., Mérő, T. O., Puky, M., Lengyel, S. (2015): Spatiotemporally
 474 variable management by grazing and burning increases marsh diversity and
 475 benefits amphibians: a field experiment. *Biol. Cons.* **192**: 237-246.

476 Mizsei, E., Jablonski, D., Végvári, Z., Lengyel, S., Szabolcs, M. (in press). Distribution
 477 and diversity of reptiles in Albania: a novel database in a Mediterranean hotspot.
 478 *Amphibia-Reptilia*.

479 Mizsei, E., Üveges, B., Vági, B., Szabolcs, M., Lengyel, S., Pfiögler, W.P., Nagy, Z.T.,
 480 Tóth J.P. (2016): Species distribution modelling leads to the discovery of new
 481 populations of one of the least known European snakes, *Vipera ursinii graeca*, in
 482 Albania. *Amphibia-Reptilia* **37**: 55-68.

483 Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Kent, J. (2000):
 484 Biodiversity hotspots for conservation priorities. *Nature* **403** (24): 853-858.

485 Ord, J.K., Getis A. (1995): Local spatial autocorrelation statistics: distribution issues
 486 and an application. *Geogr. Anal.* **27** (4): 286-306.

487 Oruçi, S. (2010): Data on geographical distribution and habitats of the *Rana epirotica*
 488 in Albania. *Natura Montenegrina* **7** (3): 419-423.

489 Özdemir, N., Gül, S., Poyarkov Jr., N. A., Kutrup, B., Tosunoğlu, Doglio, S. (2014):
 490 Molecular systematics and phylogeography of *Bufo variabilis* (syn.
 491 *Pseudepidalea variabilis*) (Pallas, 1769) in Turkey. *Turk. J. Zool.* **38**: 412-420.
 492 Pabijan, M., Zieliński, P., Dudek, K., Chloupek, M., Sotiropoulos, K., Liana, M., Babik,
 493 W. (2015): The dissection of a Pleistocene refugium: phylogeography of the
 494 smooth newt, *Lissotriton vulgaris* in the Balkans. *J. Biogeogr.* **42**: 671-683.
 495 Pabijan, M., Zieliński, P., Dudek, K., Stuglik, M., Babik, W. (2016): Isolation and gene
 496 flow in a speciation continuum in newts. *BioRxiv*.
 497 Pinheiro, J., Bates, D. (2000): *Mixed-effects models in S and S-PLUS*. New York,
 498 Springer.
 499 Pimm, S.L., Jenkins, C. (2005): Sustaining the variety of life. *Sci. Am.* **293** (3): 66-73.
 500 Ragghianti, M., Bucci, S., Casola, C., Marracci, S., Mancino, G. (2004): Molecular
 501 investigations on western Palearctic water frogs. *Ital. J. Zool.* **2**: 17-23.
 502 Ragghianti, M., Bucci, S., Guerrini, F., Mancino, G. (1999): Characterization of two
 503 repetitive DNA families (RrS1 and Rana/Pol III) in the genomes of Palearctic
 504 green water frogs. *Ital. J. Zool.* **66**: 255-263.
 505 Recuero, E., Canestrelli, D., Vörös, J., Szabó, K., Poyarkov, N. A., Arntzen, J. W.,
 506 Crnobrnja-Isailović, J., Kidov, A. A., Cogălniceanu, D., Caputo, F. P., Nascetti,
 507 G., Martínez-Solano, I. (2012): Multilocus species tree analyses resolve the
 508 radiation of the widespread *Bufo bufo* species group (Anura, Bufonidae). *Mol.*
 509 *Phylogenet. Evol.* **62**: 71-86.
 510 Rocchini, D., Hortal, J., Lengyel, S., Lobo, J.M., Jimenez-Valverde, A., Ricotta, C.,
 511 Bacaro, G., Chiarucci, A. (2011): Accounting for uncertainty when mapping

species distributions: the need for maps of ignorance. *Prog. Phys. Geogr.* **35**:
211-226.

Schneider, H., Haxhiu, I. (1994): Mating-call analysis and taxonomy of the water frogs
in Albania (Anura: Ranidae). *Zool. Jahrb. Syst.* **121**: 248-262.

Scribner, K.T., Arntzen, J.W., Cruddace, N., Oldham, R.S., Burke, T. (2001):
Environmental correlates of toad abundance and population genetic diversity.
Biol. Cons. **98**: 201-210.

Shehu, M., Serravalle, F., Alfonso, G., Moscatello, S., Belmonte, G. (2009): The alpine
lake Gistova (Mount Gramos, Albania-Greece border) biodiversity of an isolated
microcosm. *Thalassia Sal.* **32**: 53-62.

Shkurti, R. (2013): The assessment of biological and economic capacity for the park of
Viroi, Gjirokastra, Albania. The 1st International Conference on Research and
Education – Challenges Towards the Future, Shkodra, Albania.

Sillero, N., Campos, J., Bonardi, A., Corti, C., Creemers, R., Crochet, P-A., Crnobrnja-
Isailović, J., Denoël, M., Ficetola, G.F., Gonçalves, J., Kuzmin, S., Lymberakis,
P., de Pous, P., Rodríguez, A., Sindaco, R., Speybroek, J., Toxopeus, B., Vieites,
D.R., Vences, M., (2014a): Updated distribution and biogeography of
amphibians and reptiles of Europe. *Amphibia-Reptilia* **35**: 1-31.

Sillero, N., Oliviera, M. A., Sousa, P., Gonçalves-Seco, L. (2014b): Distributed
database system of the New Atlas of Amphibians and Reptiles in Europe: the
NA2RE project. *Amphibia-Reptilia* **35**: 33-39.

Sodhi, N. S., Bickford, D., Diesmos, A. C., Lee, T. M., Koh, L. P., Brook, B. W.,
Sekercioglu, C. H., Bradshaw, C. J. A. (2008): Measuring the meltdown: drivers
of global amphibian extinction and decline. *Plos One* **3** (2): e1636

536 Speybroeck, J., Beukema, W., Bok, B., Van Der Voort, J., Velikov, I. (2016): Field
537 guide to the amphibians and reptiles of Britain and Europe. London,
538 Bloomsbury.

539 Stuart, S. N., Chanson, J. S., Cox, N. A., Young, B. E., Rodrigues, A. N. S. L.,
540 Fischman, D. L., Waller, R. W. (2004): Status and trends of amphibian declines
541 and extinctions worldwide. *Science* **306**: 1783-1786.

542 Szabolcs, M., Mizsei, E. (in press): First record of the eastern spadefoot toad (*Pelobates*
543 *syriacus* Boettger, 1889) in Albania. *North-West. J. Zool.*

544 Uhrin, M., Šíbl, J. (1996): Další faunistické poznámky z Albánska – pp. 5-6 in:
545 Anonymus (eds): Sborník abstrakt III. konference České herpetologické
546 společnosti. Česká herpetologická společnost, Mariánské Lázně, 11 pp.

547 Uzzell, T., Crnobrnja-Isailović, J. (2009): *Pelophylax shqipericus*. The IUCN Red List
548 of Threatened Species.e.T58715A11829016.

549 Uzzell, T., Lymberakis, P., Haxhiu, I. (2009): *Pelophylax epeiroticus*. The IUCN Red
550 List of Threatened Species.e.T58592A11793456.

551 Vági, B., Kovács, T., Bancilă, R., Hartel, T., Anthony, B. P. (2013): A landscape-level
552 study on the breeding site characteristics of ten amphibian species in Central
553 Europe. *Amphibia-Reptilia* **34**: 63-73.

554 Van Buskirk, J., (2005): Local and landscape influence on amphibian occurrence and
555 abundance. *Ecology* **86**: 1936-1947.

556 Vojar, J., Havlíková, B., Solský, M., Jablonski, D., Iković, V., Baláž, V. (2017):
557 Distribution, prevalence and amphibian hosts of *Batrachochytrium*
558 *dendrobatidis* in the Balkans. *Salamandra* **53** (1): 44-49.

559 Werner, F. (1920): Zur Kenntnis der Reptilien- und Amphibienfauna Albaniens. Zool.
560 Anz. **17** (1/2).

561 Table 1.Environmental variables used in this study.

Predictor	Description	Data source
BIO PC1	“Temperature” principal component BIO1 = Annual Mean Temperature BIO6 = Min Temperature of Coldest Month BIO11 = Mean Temperature of Coldest Quarter	Hijmans et al. 2005
BIO PC2	“Precipitation” principal component BIO12 = Annual Precipitation BIO16 = Precipitation of Wettest Quarter BIO19 = Precipitation of Coldest Quarter	Hijmans et al. 2005
BIO PC3	“Temperature variation” principal component BIO2 = Mean Diurnal Range (Mean of monthly (max temp - min temp)) BIO4 = Temperature Seasonality (standard deviation *100) BIO7 = Temperature Annual Range (BIO5-BIO6)	Hijmans et al. 2005
BIO PC4	“Precipitation variation” principal component BIO9 = Mean Temperature of Driest Quarter BIO10 = Mean Temperature of Warmest Quarter BIO15 = Precipitation Seasonality (Coefficient of Variation)	Hijmans et al. 2005
CORINE DIV	Shannon diversity of CORINE Land cover in 10×10 km cells	European Environment Agency
ALT MEAN	Mean of altitude values in 10×10 km cells, calculated from the SRTM near 90 m data	CGIA-CSI
ALT SD	Standard deviation of altitude values in 10×10 km cells, calculated from the SRTM near 90 m data	CGIA-CSI
SEA DIST	Min distance of 10×10 km cells centroids from the Adriatic sea coast	present study

563 Table 2. List of amphibian species in Albania with their number of records, Extent of Occurrence (EOO) and Distribution type.

Species	Total records	Published records	Unpublished records	N of presence 10×10 km cells	EOO (km ²)	Distribution type
<i>Bombina variegata</i>	136	46	90	67	24457	Southern-European
<i>Bufo bufo</i>	70	27	43	50	25977	European
<i>Bufo viridis/variabilis</i>	96	36	60	53	25945	Turano-European-Mediterranean
<i>Hyla arborea</i>	48	26	22	33	21559	European-Mediterranean
<i>Ichthyosaura alpestris</i>	63	43	20	31	12047	European
<i>Lissotriton graecus</i>	55	34	21	47	25065	European
<i>Pelobates syriacus</i>	3	3	0	1	1	Eastern-Mediterranean
<i>Pelophylax</i> spp.	399	221	178	181	26010	Eurasian
<i>Pelophylax epeiroticus</i>	8	5	3	6	597	Eastern-Mediterranean
<i>Pelophylax ridibundus</i>	59	54	5	41	23735	Turano-European
<i>Pelophylax shqipericus</i>	25	21	5	19	7028	Eastern-Mediterranean
<i>Rana dalmatina</i>	54	28	27	35	22938	Southern-European
<i>Rana graeca</i>	69	16	53	43	23250	Eastern-Mediterranean
<i>Rana temporaria</i>	16	15	1	14	6536	European
<i>Salamandra atra</i>	6	2	4	3	335	Central-European
<i>Salamandra salamandra</i>	42	31	11	40	24424	European-Mediterranean
<i>Triturus macedonicus</i>	39	14	25	29	21279	Eastern-Mediterranean
Total	1097	539	558	238		

565 Table 3. Predictor importance in the two GLMM models.

Presence		Shannon diversity	
Predictor	Importance	Predictor	Importance
CORINE DIV	1.000	CORINE DIV	1.000
BIO PC2	0.903	BIO PC2	0.427
BIO PC3	0.756	BIO PC4	0.291
BIO PC1	0.217	ALT SD	0.181
SEA DIST	0.204	ALT MEAN	0.087
ALT SD	0.122	BIO PC3	0.069
ALT MEAN	0.095	SEA DIST	0.000
BIO PC4	0.000	BIO PC1	0.000

566 Table 4. Parameter estimates and AIC values of the best ($\Delta AIC_c < 2$) GLMM models fitted on the presence and diversity of amphibians in
567 Albania.

Variable	Model	CORINE DIV	BIO PC2	BIO PC3	BIO PC1	SEA DIST	ALT SD	ALT MEAN	BIO PC4	df	AICc	ΔAIC_c
Presence	1	1.37504	-0.08859	-0.10412						8	2890.266	0.000
	2	1.29567	-0.08835							7	2891.487	1.222
	3	1.38543	-0.09603	-0.10198			0.00052			9	2891.656	1.391
	4	1.37858	-0.14665		0.08909	-0.00001				9	2891.834	1.568
	5	1.38769	-0.08935	-0.10471	0.01423					9	2891.945	1.679
	6	1.35787		-0.10515						7	2892.121	1.856
	7	1.38320	-0.08723	-0.10331				0.00006		9	2892.161	1.896
	8	1.37949	-0.08434	-0.11082		0.00000				9	2892.204	1.939
Diversity	1	0.31683								4	632.902	0.000
	2	0.31918	-0.02937							5	633.056	0.154
	3	0.32254							-0.04277	5	633.357	0.455
	4	0.32408	-0.03133						-0.04564	6	633.641	0.739
	5	0.29448	-0.03213				0.00037			6	634.041	1.139
	6	0.32376						0.00008		5	634.095	1.193
	7	0.29240					0.00032			5	634.563	1.661
	8	0.33340		-0.02531						5	634.663	1.761

568 Table 5. Model averaged parameter estimates of GLMM fitted on amphibian presence and
 569 MCMCglmm fitted on Shannon diversity of amphibians. Significant parameter estimates are
 570 indicated in bold.

Response	Main effect	Estimate	S.E.	z value	P
Presence	(Intercept)	-5.172	0.804	6.428	0.000
	CORINE DIV	1.367	0.227	6.015	0.000
	BIO PC2	-0.096	0.046	2.093	0.036
	BIO PC3	-0.105	0.053	1.982	0.047
	BIO PC1	0.053	0.049	1.072	0.284
	SEA DIST	0.000	0.000	0.766	0.444
	ALT SD	0.001	0.000	1.109	0.267
	ALT MEAN	0.000	0.000	0.326	0.745
	BIO PC4	0.000	0.000	-0.191	0.848
Response	Main effect	Estimate	Lower 95% CI	Upper 95% CI	P
Diversity	(Intercept)	0.224	-0.636	1.001	0.558
	CORINE DIV	0.325	0.175	0.495	0.001
	BIO PC2	-0.048	-0.101	0.004	0.078
	BIO PC4	-0.056	-0.145	0.028	0.228
	ALT SD	0.000	-0.001	0.001	0.772
	ALT MEAN	0.000	-0.001	0.000	0.606
	BIO PC3	-0.001	-0.069	0.063	0.962
	SEA DIST	0.000	0.000	0.000	0.474
	BIO PC1	0.037	-0.046	0.108	0.302

572 Figure 1. Geographic map of the study area indicating toponymics mentioned in the text.

573 Figure 2. Number of records by year of publication (published sources) or year of data
574 collection (unpublished sources). Vertical line indicates the year when the former isolationist
575 political system ended in Albania (1991).

576


577 Figure 3. Sources of occurrence records of amphibian species used in the present study (A),
578 sampling hotspots (GiZ score > 1.0) and coldspots (GiZ score < -1.0) (B), and amphibian
579 species richness (numbers) and Shannon diversity (shading) (C) in Albania on a 10×10 km
580 grid.

581


582 Figure 4. Altitudinal distribution of amphibian species and frequency of occurrence records
583 by altitude in Albania. Box-and-whiskers plots show the median (horizontal line), the 25th
584 and 75th percentile (bottom and top of box, respectively), minimum and maximum values
585 (lower and upper whiskers, respectively) and outliers (circles). Grey line (red in the colour
586 version) is the frequency distribution of altitudinal values in Albania.


587


588 Figure 5. Species presence and Shannon diversity as a function of the most important
589 predictors identified by GLMM model selection (for abbreviations, see Table 1).




















Figure4col







[Click here to access/download](#)

Supplementary File

ALFA_amphibians_AMRE_supplementary.docx

